

Appendices

Glossary

AGGRADATION: The geologic process in which streambeds, floodplains, and the bottoms of other water bodies are raised in elevation by the deposition of material eroded and transported from other areas. It is the opposite of degradation.

ALEVIN: The life stages of salmonids that occurs after eggs have hatched but before young emerge from the gravel nests where they have incubated. Alevin still have yolk sacs attached to provide them with nutrition within the nest.

ALLUVIUM: A general term for all deposits resulting directly or indirectly from the sediment transport of streams, thus including the sediments lay down in riverbeds, floodplains, lakes, fans, and estuaries.
ALLUVIAL adj.

ANADROMOUS: Fish that leave freshwater and migrate to the ocean to mature then return to freshwater to spawn. Salmon, steelhead, and shad are examples.

ANTHROPOGENIC: Caused by humans.

ARCINFO: ESRI (Environmental Systems Research Institute) proprietary software, which provides a complete GIS data creation, update, query, mapping, and analysis system.

AERIAL: Having to do with or done by aircraft. For example, aircraft equipped with cameras capture images of the earth in air photos.

BANKFULL DISCHARGE: The discharge corresponding to the stage at which the floodplain of a particular stream reach begins to be flooded; the point at which bank overflow begins.

BANKFULL WIDTH: The width of the channel at the point at which overbank flooding begins.

BASIN: see watershed.

BED SUBSTRATE: The materials composing the bottom of a stream.

BENTHIC: The collection of organisms living on or in sea, river, or lake bottoms.

BOULDER: Stream substrate particle larger than 10 inches (256 millimeters) in diameter.

CALWATER: A set of standardized watershed boundaries for California nested into larger previously standardized watersheds and meeting standardized delineation criteria.

CANOPY: The overhead branches and leaves of streamside vegetation.

CANOPY COVER: The vegetation that projects over the stream.

CANOPY DENSITY: The percentage of the sky above the stream screened by the canopy of plants, sometimes expressed by species.

CENTROID: The center of water mass of a flowing stream at any location. This location usually correlates well with the thalweg, or deepest portion of the stream. Sampling in the centroid is intended to provide a reasonably representative sample of the main stream.

CHANNEL: A natural or artificial waterway of perceptible extent that periodically or continuously contains moving water. It has a definite bed and banks, which serve to confine the water.

COAST RANGE: A string of mountain ranges along the Pacific Coast of North America from Southeastern Alaska to lower California.

COBBLE: Stream substrate particles between 2.5 and 10 inches (64 and 256 millimeters) in diameter.

COLLUVIUM: A general term for loose deposits of soil and rock moved by gravity; e.g. talus.

CONIFEROUS: Any of various mostly needle-leaved or scale-leaved, chiefly evergreen, cone-bearing gymnospermous trees or shrubs such as pines, spruces, and firs.

CONSUMPTIVE USE OF WATER: Occurs when water is taken from a stream and not returned.

COVER: Anything that provides protection from predators or ameliorates adverse conditions of streamflow and/or seasonal changes in metabolic costs. May be instream cover, turbulence, and/or overhead cover, and may be for the purpose of escape, feeding, hiding, or resting.

DEBRIS: Material scattered about or accumulated by either natural processes or human influences.

DEBRIS JAM: Logjam. Accumulation of logs and other organic debris.

DEBRIS LOADING: The quantity of debris located within a specific reach of stream channel, due to natural processes or human activities.

DECIDUOUS: A plant (usually a tree or shrub) that sheds its leaves at the end of the growing season.

DEGRADATION: The geologic process in which stream beds and floodplains are lowered in elevation by the removal of material. It is the opposite of aggradation.

DEMOGRAPHY: The study of the characteristics of populations, such as size, growth, density, distribution, and vital statistics.

DEPOSITION: The settlement or accumulation of material out of the water column and onto the streambed. Occurs when the energy of flowing water is unable to support the load of suspended sediment.

DEPTH: The vertical distance from the water surface to the streambed.

DISCHARGE: Volume of water flowing in a given stream at a given place and within a given period of time, usually expressed as cubic meters per second (m³/sec), or cubic feet per second (cfs).

DISSOLVED OXYGEN (DO): The concentration of oxygen dissolved in water, expressed in mg/l or as percent saturation, where saturation is the maximum amount of oxygen that can theoretically be dissolved in water at a given altitude and temperature.

DIVERSION: A temporal removal of surface flow from the channel.

ECOTONE: A transition area between two distinct habitats that contains species from each area, as well as organisms unique to it.

EMBEDDEDNESS: The degree that larger particles (boulders, rubble, or gravel) are surrounded or covered by fine sediment. Usually measured in classes according to percentage of coverage of larger particles by fine sediments.

ECOLOGICAL MANAGEMENT DECISION SUPPORT (EMDS): An application framework for knowledge-based decision support of ecological landscape analysis at any geographic scale.

EMBRYO: An organism in its early stages of development, especially before it has reached a distinctively recognizable form.

ENDANGERED SPECIES: Any species which is in danger of extinction throughout all or a significant portion of its range other than a species of the Class Insecta determined by the Secretary to constitute a pest whose protection under the provisions of this Act would present an overwhelming and overriding risk to man.

EROSION: The group of natural processes, including weathering, dissolution, abrasion, corrosion, and transportation, by which material is worn away from the earth's surface. *EROSIONAL adj.*

ESTUARY: A water passage where the tide meets a river current.

EXTIRPATION: To destroy totally; exterminate.

EXTINCTION: The death of an entire species.

FILL: A) The localized deposition of material eroded and transported from other areas, resulting in a change in the bed elevation. This is the opposite of scour; B) The deliberate placement of (generally) inorganic materials in a stream, usually along the bank.

FINE SEDIMENT: The fine-grained particles in stream banks and substrate. Those are defined by diameter, varying downward from 0.24 inch (6 millimeters).

FISH HABITAT: The aquatic environment and the immediately surrounding terrestrial environment that, combined, afford the necessary biological and physical support systems required by fish species during various life history stages.

FLATWATERS: In relation to a stream, low velocity pool or run habitat.

FLOOD: Any flow that exceeds the bankfull capacity of a stream or channel and flows out of the floodplain; greater than bankfull discharge.

FLOODPLAIN: The area bordering a stream over which water spreads when the stream overflows its banks at flood stages.

FLOW: A) The movement of a stream of water and/or other mobile substances from place to place; B) the movement of water, and the moving water itself; C) the volume of water passing a given point per unit of time. Discharge.

FLUVIAL: Relating to or produced by a river or the action of a river. Situated in or near a river or stream.

FRESHETS: A sudden rise or overflowing of a small stream as a result of heavy rains or rapidly melting snow.

GENETIC DRIFT: The random change of the occurrence of a particular gene in a population.

GEOGRAPHIC INFORMATION SYSTEM (GIS): A collection of computer hardware, software, and geographic data used for capturing, managing, analyzing, and displaying all forms of geographically referenced information on the Earth's surface. Typically, a GIS is used to produce graphics (maps) on the screen or on paper to convey the results of analyses.

GEOMORPHOLOGY: The study of surface forms on the earth and the processes by which these develop.

GRADIENT: The slope of a streambed or hillside. For streams, gradient is quantified as the vertical distance of descent over the horizontal distance the stream travels.

GRAVEL: Substrate particle size between 0.08 and 2.5 inches (2 and 64 millimeters) in diameter.

GULLY: A deep ditch or channel cut in the earth by running water after a prolonged downpour.

HABITAT: The place where a population lives and its surroundings, both living and nonliving; includes the provision of life requirements such as food and shelter.

HABITAT CONSERVATION PLAN: A document that describes how an agency or landowner will manage their activities to reduce effects on vulnerable species. An HCP discusses the applicant's proposed activities and describes the steps that will be taken to avoid, minimize, or mitigate the take of species that are covered by the plan.

HABITAT TYPE: A land or aquatic unit, consisting of an aggregation of habitats having equivalent structure, function, and responses to disturbance.

HETEROZYGOSITY: The presence of different alleles at one or more loci on homologous chromosomes.

HIERARCHY: A series of ordered groupings of people or things within a system.

HYDROGRAPH: A graph showing, for a given point on a stream, the discharge, stage, velocity, or other property of water with respect to time.

HYDROLOGY: The science of water, its properties, phenomena, and distribution over the earth's surface.

HYDROGRAPHIC UNIT: A watershed designation at the level below Hydrologic Region and above Hydrologic Sub-Area.

HYPOTHESIS: A tentative explanation for an observation, phenomenon, or scientific problem that can be tested by further investigation.

INBREEDING: The breeding of related individuals within an isolated or a closed group of organisms.

INBREEDING DEPRESSION: The exposure of individuals in a population to the effects of deleterious recessive genes through matings between close relatives.

INCUBATION: Maintaining something at the most favorable temperature for its development.

INSTREAM COVER: Areas of shelter in a stream channel that provide aquatic organisms protection from predators or competitors and/or a place in which to rest and conserve energy due to a reduction in the force of the current.

INTERMITTENT STREAM: A stream in contact with the ground water table that flows only at certain times of the year when the ground water table is high and/or when it receives water from springs or from some surface source such as melting snow in mountainous areas. It ceases to flow above the streambed when losses from evaporation or seepage exceed the available stream flow. Seasonal.

KNOWLEDGE BASE: An organized body of knowledge that provides a formal logical specification for the interpretation of information.

LAGOON: A shallow body of water, especially one separated from a sea by sandbars or coral reefs.

LIMITING FACTOR: Environmental factor that limits the growth or activities of an organism or that restricts the size of a population or its geographical range.

LARGE WOODY DEBRIS (LWD): A large piece of relatively stable woody material having a diameter greater than 12 inches (30 centimeters) and a length greater than 6 feet (2 meters) that intrudes into the stream channel. Large organic debris.

MACROINVERTEBRATE: An invertebrate animal (animal without a backbone) large enough to be seen without magnification.

MAINSTEM: The principal, largest, or dominating stream or channel of any given area or drainage system.

MELANGE: A mappable body of rock that includes fragments and blocks of all sizes, both exotic and native, embedded in a fragmented and generally sheared matrix.

MIGRATION: The periodic passage from one region to another for feeding or breeding.

NETWEAVER: A knowledge-based development system. A meta database that provides a specification for interpreting information.

NUTRIENT: A nourishing substance; food. The term *nutrient* is loosely used to describe a compound that is necessary for metabolism.

ONCORHYNCHUS: A genus of the family salmonidae (salmons and trouts). They are named for their hooked (onco) nose (rhynchus).

ORGANIC DEBRIS: Debris consisting of plant or animal material.

ORTHOPHOTOQUADS: A combined aerial photo and planimetric quad map (with no indication of contour) without image displacements and distortions.

PERMANENT STREAM: A stream that flows continuously throughout the year. Perennial.

pH: A measure of the hydrogen ion activity in a solution, expressed as the negative log₁₀ of hydrogen ion concentration on a scale of 0 (highly acidic) to 14 (highly basic) with a pH of 7 being neutral.

PLATE TECTONICS: A theory in which the earth's crust is divided into mobile plates which are in constant motion causing earthquake faults, volcanic eruptions, and uplift of mountain ranges.

PHOTOGRAMMETRY: Photogrammetry is the technique of measuring objects (2D or 3D) from photographs or other digital imagery. May also be referred to as Remote Sensing.

PRODUCTIVITY: A) Rate of new tissue formation or energy utilization by one or more organisms; B) Capacity or ability of an environmental unit to produce organic material; C) The ability of a population to recruit new members by reproduction.

REDD: A spawning nest made by a fish, especially a salmon or trout.

REFERENCE CONDITIONS: Minimally impaired conditions that provide an estimate of natural variability in biological condition and habitat quality.

- RIFFLE:** A shallow area extending across a streambed, over which water rushes quickly and is broken into waves by obstructions under the water.
- RILL:** An erosion channel that typically forms where rainfall and surface runoff is concentrated on slopes. If the channel is larger than one square foot in size, it is called a gully.
- RIPARIAN:** Pertaining to anything connected with or immediately adjacent to the banks of a stream or other body of water.
- RIPARIAN AREA:** The area between a stream or other body of water and the adjacent upland identified by soil characteristics and distinctive vegetation. It includes wetlands and those portions of floodplains and valley bottoms that support riparian vegetation.
- RIPARIAN VEGETATION:** Vegetation growing on or near the banks of a stream or other body of water on soils that exhibit some wetness characteristics during some portion of the growing season.
- RUBBLE:** Stream substrate particles between 2.5 and 10 inches (64 and 256 millimeters) in diameter.
- SALMONID:** Fish of the family *Salmonidae*, including salmon, trout, chars, whitefish, ciscoes, and graylings.
- SCOUR:** The localized removal of material from the streambed by flowing water. This is the opposite of fill.
- SEDIMENT:** Fragmented material that originates from weathering of rocks and decomposition of organic material that is transported by, suspended in, and eventually deposited by water or air, or is accumulated in beds by other natural phenomena.
- SERIAL STAGES:** The series of relatively transitory plant communities that develop during ecological succession from bare ground to the climax stage.
- SHEAR:** A deformation resulting from stresses that cause contiguous parts of a body to slide relatively to each other in a direction parallel to their plane of contact.
- SILVICULTURE:** The care and cultivation of forest trees; forestry.
- SMOLT:** Juvenile salmonid one or more years old that has undergone physiological changes to cope with a marine environment, the seaward migration stage of an anadromous salmonid.
- SMOLTIFICATION:** The physiological change adapting young anadromous salmonids for survival in saltwater.
- SPAWNING:** To produce or deposit eggs.
- STADIA RODS:** Graduated rods observed through a telescopic instrument while surveying to determine distances and elevation.
- STAGE:** The elevation of a water surface above or below an established datum or reference.
- STREAM:** (includes creeks and rivers): A body of water that flows at least periodically or intermittently through a bed or channel having banks and supports fish or other aquatic life. This includes watercourses having a surface or subsurface flow that supports or has supported riparian vegetation.
- STREAM BANK:** The portion of the channel cross section that restricts lateral movement of water at normal water levels. The bank often has a gradient steeper than 45 degrees and exhibits a distinct break in slope from the stream bottom. An obvious change in substrate may be a reliable delineation of the bank.
- STREAM CLASSIFICATION:** Various systems of grouping or identifying streams possessing similar features according to geomorphic structure (e.g. gradient, water source, spring, and creek), associated biota (e.g. trout zone), or other characteristics.
- STREAM CORRIDOR:** A stream corridor is usually defined by geomorphic formation, with the corridor occupying the continuous low profile of the valley. The corridor contains a perennial, intermittent, or ephemeral stream and adjacent vegetative fringe.
- STREAM REACH:** A section of a stream between two points.

- SUBSTRATE:** The material (silt, sand, gravel, cobble, etc.) that forms a stream or lakebed.
- SUBWATERSHED:** One of the smaller watersheds that combine to form a larger watershed.
- TAKE:** To harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct.
- TERRACE:** A former floodplain underlain by sediment deposited by a stream when the stream was flowing at a higher level; typically forming a relatively level bench along a valley side adjacent to a recent floodplain.
- TERRAIN:** A tract or region of the earth's surface considered as a physical feature, an ecological environment, or a site of some planned activity of man.
- TERRANE:** A term applied to a rock or group of rocks and to the area in which they crop out. The term is used in a general sense and does not imply a specific rock unit.
- THALWEG:** The line connecting the lowest or deepest points along a streambed.
- THREATENED SPECIES:** Any species that is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range.
- TOPOGRAPHY:** The general configuration of a land surface, including its relief and the position of its natural and man-made features.
- TRIBUTARY:** A stream feeding, joining, or flowing into a larger stream. Feeder stream, side stream.
- UNDERCUT BANK:** A bank that has had its base cut away by the water action along man-made and natural overhangs in the stream.
- VELOCITY:** The time rate of motion; the distance traveled divided by the time required to travel that distance.
- V*:** Measures of percent sediment filling of a stream pool with deposits such as silt, sand, and gravel compared to the total volume.
- WATER RIGHT:** The right to draw water from a particular source, such as a lake, irrigation canal, or stream. Often used in the plural.
- WATERSHED ASSESSMENT:** An interdisciplinary process of information collection and analysis that characterizes current watershed conditions at a course scale.
- WATERSHED:** Total land area draining to any point in a stream, as measured on a map, aerial photograph or other horizontal plane. Also called catchment area, watershed, and basin.
- WATERSHED MANAGEMENT AREA (WMA):** In the context of the North Coast Regional Water Quality Control Board's Watershed Management Initiative, this represents a grouping of smaller watersheds into a larger area for identifying and addressing water quality problems, e.g., the Humboldt WMA includes all watersheds draining to the ocean or bays north of the Eel River to and including Redwood Creek.
- WETLAND:** An area subjected to periodic inundation, usually with soil and vegetative characteristics that separate it from adjoining non-inundated areas.

List of Abbreviations

CalEPA	California Environmental Protection Agency
CCWPAP	California Coastal Watershed Planning and Assessment Program
CDF	California Department of Forestry and Fire Protection
CEMPR	Center for Manpower Resources
CEQA	California Environmental Quality Act
CESA	California Endangered Species Act
CFS	Cubic Feet per Second
CDFG	California Department of Fish and Game
DOC/CGS	California Department of Conservation-California Geological Survey
DPR	California Department of Parks and Recreation
DWR	California Department of Water Resources
EMDS	Ecological Management Decision Support
EPA	Environmental Protection Agency
ESA	Federal Endangered Species Act
ESU	Evolutionarily Significant Units
FRGP	Fishery Restoration Grants Program
FISRWG	Federal Interagency Stream Restoration Working Group
FPA	Z'Berg-Nejedly Forest Practice Act
GIS	Geographic Information System
GMA	Graham Mathews and Associates
HA	Hydrologic Area
HCP	Habitat Conservation Plan
HR	North Coast Hydrologic Region
HAS	Hydrologic Sub-area
HTC	Hawthorne Timber Company
HU	Hydrologic Unit
IA	Integrated Analysis
JDSF	Jackson State Demonstration Forest
LFA	Limiting Factor Analysis
LP	Louisiana Pacific Lumber Company
LWD	Large Woody Debris
MOU	Memorandum of Understanding
MRC	Mendocino Redwood Company, LLC
MWAT	Maximum Weekly Average Temperature
NCRWQCB	North Coast Regional Water Quality Control Board
NCWAP	North Coast Watershed Assessment Program
NEPA	National Environmental Policy Act
NMFS	National Marine Fisheries Service
PW	Planning Watershed
RM	River Mile
SONAR	Mendocino High School of Natural Resources
SRP	Scientific Review Panel
SWRCB	California State Water Resources Control Board
TMDL	Total Maximum Daily Load
TPZ	Timber Production Zone
USFS	United States Forest Service
USGS	United States Geologic Survey
WMA	Watershed Management Area
WQO	Water Quality Objectives
WRIMS	State Water Resources Control Board Water Right Information System

Bibliography

- Anonymous. 1904. Stocking Report. Mendocino Dispatch Democrat. Mendocino, California.
- Arvola, T. 1977. Regulation of logging in California, 1945-1975. Division of Forestry, State of California. Sacramento, California.
- Barber, T. 2003 (Draft). Mendocino Woodlands Erosion Prevention and Implementation Plan. For Mendocino Woodlands State Park by Ridge to River Watershed Sciences, Restoration, Forestry, and Water Quality. Westport, California.
- Barnhart, R. 1994. Salmon and steelhead populations of the Klamath-Trinity Basin, California. Proceedings of the Klamath Basin Fisheries Symposium, Humboldt State University, Arcata, California, California Cooperative Fisheries Research Unit.
- Big River Watershed Council. 1997. Watershed Guidelines for Big River Watershed, Mendocino County, California. By the Big River Watershed Council for National Marine Fisheries Service, Ukiah, California. 7p.
- Bilby, R.E. and J.W. Ward. 1989. Changes in the characteristics and function of woody debris with increasing size of streams in western Washington. Transactions of the American Fisheries Society 118:368-378.
- Bisson P.A., R.E. Bilby, M.D. Bryant, C.A. Dolloff, G.B. Grette, R.A. House, M.L. Murphy, K.V. Koski, and J.R. Sedell. 1987. Large woody debris in forested streams in the Pacific Northwest: past, present, and future. Streamside Management: Forestry and Fishery Interactions, pp. 143-190. Contribution 57. University of Washington Institute of Forest Resources, Seattle, Washington.
- Bjornn, T.C. and D. W. Reiser. 1991. Habitat Requirements of Salmonids in Streams. American Fisheries Society Special Publication 19:83-138.
- Blake, M. C. and D. Jones. 1974. Origin of Franciscan Mélanges in northern California. Modern and ancient geosynclinal sedimentation. R. D. Jr. and R.H. Shaver, Society of Economic Paleontologists and Mineralogists. Special Publication 19: 345-357.
- Blake, M. C. and D. Jones. 1981. The Franciscan Assemblage and Related Rocks in Northern California: A reinterpretation. The geotectonic development of California. W. Ernst. Prentice-Hall, Inc., Englewood Cliffs, New Jersey. Ruby Volume I: 307-328.
- Bleier, C., Downie, S., Cannata, S., Henly, R., Walker, R., Keithley, C. Scruggs, M.; Custis, K., Clements, J. and R. Klamt. 2003. North Coast Watershed Assessment Program Methods Manual. California Resources Agency and California Environmental Protection Agency, Sacramento, California.
- Brett, J.R. 1952. Temperature tolerance in young Pacific salmon, Genus *Oncorhynchus*. J. Fish. Res. Bd. Can. 9:265-323.
- Britschgi, D. E. and M. L. Marcus. 1981. Aquatic Life. Big River: the Natural History of and Endangered Northern California Estuary. S. Warrick and E. Wilcox. Environmental Field Program, University of California Santa Cruz. Santa Cruz, California. pp 47-98.
- Brown, P. and W. Baxter. 2003. "Fire History in Coastal Redwood Forests of the Mendocino Coast, California." Northwest Science 77(2): 147-158.
- Brungs, W.S. and B.R. Jones. 1977. Temperature Criteria for Freshwater Fish: Protocols and Procedures. EPA-600/3-77-061. Environ. Research Lab, Ecological Resources Service, U.S. Environmental Protection Agency, Office of Research and Development, Duluth, MN.
- Burns, J. 1971. "The Carrying Capacity for Juvenile Salmonids in Some Northern California Streams." California Fish and Game 57(1): 44-57.
- Cafferata, P. and T. Spittler. 1998. Logging Impacts of the 1970's vs. the 1990's in the Caspar Creek Watershed, USFS.

- California State Board of Forestry and Fire Protection (BOF). 1999. Hillslope monitoring program: Monitoring results from 1996 through 1998. Interim report prepared by the Monitoring Study Group (MSG). Sacramento, CA. 70 p.
- California Department of Forestry (CDF). 1947. Black and white aerial photos, Flight CDFI, Flight Line 1, Frames 5-6, 74-83, 102-107, 124-129; Flight Line 1A, Frames 89-97; Flight Line 2, Frames 19-21; Flight Line 4, Frames 158-163; Flight Line 5; Frames 9-19, 43-47; Flight Line 6, Frames 14-18; Flight Line 7, Frames 45-47, 101-114, 125-135, 159-168; Flight Line 9, Frames 12-17; scale 1:20,000.
- CDF. 1999. California Department of Forestry and Fire Protection. Draft Habitat Conservation Plan/Sustained Yield Plan for Jackson Demonstration State Forest.
- California Department of Conservation. California Geological Survey (CGS) 2004. Engineering Geologic Resource Assessment Big River State Park, Mendocino County, California: 32 p.
- California Department of Fish and Game (CDFG). 1958. Stream Survey - North Fork Big River.
- CDFG. 1957-1958. Stream Survey - South Fork Big River.
- CDFG. 1958. Stream Survey - East Branch North Fork Big River.
- CDFG. 1958. Stream Survey - James Creek.
- CDFG. 1958. Stream Survey - West Fork James Creek.
- CDFG. 1958. Stream Survey - South Fork Big River Tributary #1.
- CDFG. 1958. Stream Survey - South Fork Big River Tributary #2.
- CDFG. c. 1959. Stream Survey - Tramway Gulch.
- CDFG. 1959. Stream Survey - Mainstem Big River, California Department of Fish and Game: pp 1-5.
- CDFG. 1959. Stream Survey - Little North Fork Big River.
- CDFG. 1959. Stream Survey - Two Log Creek.
- CDFG. 1959. Stream Survey - North Fork Big River.
- CDFG. 1959. Stream Survey - Ramon Creek.
- CDFG. 1959. Stream Survey - Daugherty Creek (Part 1).
- CDFG. 1959. Stream Survey - Daugherty Creek (Part 2).
- CDFG. 1959. Stream Survey - Soda Creek.
- CDFG. 1959. Stream Survey - Johnson Creek (Tributary to Gates Creek).
- CDFG. 1959. Stream Survey - Snuffins Creek.
- CDFG. 1959. Stream Survey - Johnson Creek.
- CDFG. 1959. Stream Survey - Russell Brook.
- CDFG. 1959. Stream Survey - Pig Pen Gulch.
- CDFG. 1959. Stream Survey - Martin Creek.
- CDFG. 1959. Stream Survey - East Fork Martin Creek.
- CDFG. 1959. Stream Survey - Valentine Creek.
- CDFG. 1959. Stream Survey - Rice Creek.
- CDFG. 1965. California Fish and Wildlife Plan, Volume III, Supporting Data Part B Inventory (Salmon-Steelhead and Marine Resources). California Department of Fish and Game, Sacramento, California. pp 323-967.
- CDFG. 1966. Stream Survey - Two Log Creek.
- CDFG. 1966. Stream Survey - East Branch North Fork Big River.

- CDFG. 1966. Stream Survey - South Fork Big River.
- CDFG. 1966. Stream Survey - Snuffins Creek.
- CDFG. 1966. Stream Survey - Johnson Creek.
- CDFG. 1988. Electrofishing Report - Hatch Gulch.
- CDFG. 1988. Electrofishing Report - Daugherty Creek.
- CDFG. 1993. Stream Inventory Report - Daugherty Creek.
- CDFG. 1993. Stream Inventory Report - Gates Creek.
- CDFG. 1993. Stream Inventory Report - Snuffins Creek.
- CDFG. 1993. Stream Inventory Report - Johnson Creek (Tributary to Gates Creek).
- CDFG. 1995. Stream Inventory Report - Little North Fork Big River.
- CDFG. 1995. Stream Inventory Report - Soda Creek.
- CDFG. 1996. Stream Inventory Report - James Creek.
- CDFG. 1996. Stream Inventory Report - Railroad Gulch.
- CDFG. 1996. Stream Inventory Report - North Fork Big River.
- CDFG. 1997. Stream Inventory Report - Berry Gulch.
- CDFG. 1997. Stream Inventory Report - Chamberlain Creek.
- CDFG. 1997. Stream Inventory Report - North Fork Big River.
- CDFG. 1997. Stream Inventory Report - North Fork James Creek.
- CDFG. 1997. Stream Inventory Report - Rocky Gulch.
- CDFG. 1997. Stream Inventory Report - Soda Gulch.
- CDFG. 1997. Stream Inventory Report - Thompson Gulch.
- CDFG. 1997. Stream Inventory Report - Water Gulch.
- CDFG. 1997. Stream Inventory Report - Manly Gulch.
- CDFG. 1997. Stream Inventory Report - Berry Gulch Tributary.
- CDFG. 1997. Stream Inventory Report - Two Log Creek.
- CDFG. 1997. Stream Inventory Report - Water Gulch Tributary.
- CDFG. 1997. Stream Inventory Report - Park Gulch.
- CDFG. 1997. Stream Inventory Report - West Chamberlain Creek.
- CDFG. 1997. Stream Inventory Report - Gulch Sixteen.
- CDFG. 1997. Stream Inventory Report - Gulch Sixteen Tributary.
- CDFG. 1997. Stream Inventory Report - Arvola Gulch.
- CDFG. 1997. Stream Inventory Report - Lost Lake Creek.
- CDFG. 1998. Stream Inventory Report - East Branch North Fork Big River.
- CDFG. 1998. Stream Inventory Report - Two Log Creek.
- CDFG. 1998. Stream Inventory Report - Two Log Creek Tributary (Ayn Creek).
- CDFG. 1998. Stream Inventory Report - Two Log Creek Tributary (Sauerkraut Creek).
- CDFG. 2001. California's Living Marine Resources: A Status Report, University of California. 592 p.
- CDFG. 2002. Stream Inventory Report - Anderson Gulch.

- CDFG. 2002. Stream Inventory Report - Big River Tramway Gulch to North Fork Big River.
- CDFG. 2002. Stream Inventory Report - Big River from Wheel Gulch to Blind Gulch.
- CDFG. 2002. Stream Inventory Report - Biggs Gulch.
- CDFG. 2002. Stream Inventory Report - Dark Gulch.
- CDFG. 2002. Stream Inventory Report - Boardman Gulch.
- CDFG. 2002. Stream Inventory Report - Daugherty Creek.
- CDFG. 2002. Stream Inventory Report - East Branch Little North Fork Big River.
- CDFG. 2002. Stream Inventory Report - Gates Creek.
- CDFG. 2002. Stream Inventory Report - Horse Thief Creek.
- CDFG. 2002. Stream Inventory Report - Johnson Creek.
- CDFG. 2002. Stream Inventory Report - Kidwell Gulch.
- CDFG. 2002. Stream Inventory Report - Martin Creek.
- CDFG. 2002. Stream Inventory Report - Martin Creek Left Bank Tributary.
- CDFG. 2002. Stream Inventory Report - Martin Creek Right Bank Tributary #1.
- CDFG. 2002. Stream Inventory Report - Martin Creek Right Bank Tributary #2.
- CDFG. 2002. Stream Inventory Report - Mettick Creek.
- CDFG. 2002. Stream Inventory Report - Montgomery Creek.
- CDFG. 2002. Stream Inventory Report - Ramon Creek.
- CDFG. 2002. Stream Inventory Report - North Fork Ramon Creek.
- CDFG. 2002. "Stream Inventory Report - Rice Creek."
- CDFG. 2002. Stream Inventory Report - Russell Brook.
- CDFG. 2002. Stream Inventory Report - South Fork Big River.
- CDFG. 2002. Stream Inventory Report - Snuffins Creek.
- CDFG. 2002. Stream Inventory Report - Soda Creek.
- CDFG. 2002. Stream Inventory Report - Two Log Creek.
- CDFG. 2002. Stream Inventory Report - South Fork Big River Tributary #1.
- CDFG. 2002. Stream Inventory Report - South Fork Big River Tributary #2.
- CDFG. 2002. Stream Inventory Report - Valentine Creek.
- CDFG. 2002. Stream Inventory Report - Halfway House Gulch.
- CDFG. 2002. Stream Inventory Report - Johnson Creek (Tributary to Gates Creek).
- CDFG. 2002. Status review of California coho salmon north of San Francisco.
- California Department of Parks and Recreation (DPR). 2005a. Big River Watershed Restoration Proposition 40 River Parkways Grant - Preliminary Draft.
- DPR. 2005b. Big River Preliminary Plan: Resource Assessment and Recommendations. Big River Project Area, Mendocino County, California: 177 p.
- California Department of Water Resources (DWR). 1965. North Coastal Area Investigation, Appendix C, Fish and Wildlife, Dept. of Water Resources Bulletin No. 136. Dept. of Water Resources, Redding CA.
- Carver, G. A., A.S. Jayco, D.W. Valentine, W.H. Li. 1994. Coastal uplift associated with the 1992 Cape Mendocino earthquake, Northern California. *Geology* 22: 195-198.

- Center for Education and Manpower Resources (CEMR). 1979. Stream Survey - Railroad Gulch.
- Center for Education and Manpower Resources (CEMR). 1979. Stream Survey - Little North Fork Big River.
- Center for Education and Manpower Resources (CEMR). 1979. Stream Survey - Thompson Gulch.
- Center for Education and Manpower Resources (CEMR). 1979. Stream Survey - East Branch North Fork Big River.
- Cherr and J. Griffin. 1979. Department of Fish and Game Stream Inventory - Mendocino County, California. California Department of Fish and Game: 241p.
- Clapsadle, J., N. Sharples, et al. 2001. Big River Estuary Transect Fish Count. School of Natural Resources, Mendocino High School, Mendocino, California. 7p.
- Dana, G. 1978. Coastal Wetland Survey - Big River, California Department of Fish and Game: 12p.
- Downie, Scott T., C.M. LeDoux-Bloom, K. Spivak, and F. Yee, (Multi-disciplinary team leads). 2003. Albion River Basin Assessment Report. North Coast Watershed Assessment Program. California Resources Agency, and California Environmental Protection Agency, Sacramento, California.
- Durham, J. B. 1979. Geologic map of the Fort Bragg 15' quadrangle, Department of Forestry, Region 1 Geology Group.
- Durham, J. B. 1979. Geologic map of the Comptche 15' quadrangle, Department of Forestry, Region 1 Geology Group.
- Durham, J. B. 1979. Geologic map of the Willits 15' quadrangle, Department of Forestry, Region 1 Geology Group.
- Durham, J. B. 1979. Geologic map of the Navarro 15' quadrangle, Department of Forestry, Region 1 Geology Group.
- Durham, J. B. 1979. Geologic map of the Booneville 15' quadrangle, Department of Forestry, Region 1 Geology Group.
- Eilperin, J. 2006. Conservationists vie to buy forest habitat. The Washington Post March 20, 2006.
- Elwell, R. 1958. Big River Drainage Survey, Mendocino County, California. California Department of Fish and Game Intraoffice Correspondence.
- Emig, J. 1996. Barrier Debris Jam on the North Fork Big River, California Department of Fish and Game Correspondence.
- English, S. 1979. Final Draft Environmental Assessment for Protection of Big River Estuary, Mendocino County, California. US Fish and Wildlife Service, Sacramento, California..
- Escola, E. 2001-2002. Retired District Ranger, California Department of Forestry and Fire Protection.
- Evans, W. 1955. Meeting at Fort Bragg, July 19, 1955. Regarding King Salmon Project in Big River, Mendocino County, California. California Department of Fish and Game Intraoffice Correspondence.
- Federal Interagency Stream Restoration Working Group (FISRWG). 1998. Stream Corridor Restoration: Principles, Processes, and Practices. GPO Item No. 0120-A; SuDocs No. A57.6/2:EN 3/PT.653.
- Flosi, G., S. Downie, J. Hopelain, M. Bird, R. Coey, and B. Collins. 1998. California salmonid stream habitat restoration manual California Department of Fish and Game, Sacramento, California.
- Forest Ecosystem Management Assessment Team (FEMAT). 1993. Forest ecosystem management: an ecological, economic, and social assessment. U.S. Department of Agriculture, U.S. Department of Interior, U.S. Department of Commerce, U.S. Environmental Protection Agency, Portland, OR.
- Fox, K. 1983. Tectonic setting of the late Miocene, Pliocene, and Pleistocene rocks in part of the Coast Ranges north of San Francisco, California, U.S. Geological Survey: 33p.
- Frissell, C., P. Morrison, et al. 2000. Identifying Priority Areas for Salmon Conservation in the Puget Sound Basin, Pacific Biodiversity Institute.

- Fritz, E. 1942. Big River Basin as a Prospective State Forest Unit with Particular Reference to the Lands of the Mendocino Lumber Company. Berkeley, California: 23p.
- Frissell CA. 1993. Topology of extinction and endangerment of native fishes in the Pacific Northwest and California (U.S.A.). *Conservation Biology* 7: 342-354.
- Geniella, M. 1999. EPA Study Blames Rural Roads for Eel River Sediment, Natural Events Also Hurt Fish. *Press Democrat*. Santa Rosa, California.
- Geniella, M. 2006. \$2.4 Million Deal Doubles Size of Redwood Reserve. *Press Democrat*, Santa Rosa, California.
- Georgia-Pacific. 1995. Stream Inventory Report - Little North Fork Big River.
- Georgia-Pacific. 1996. Stream Inventory Report - Mainstem Big River, Georgia Pacific: pp1-8. Mainstem Big River from end of tidal zone, near confluence of Laguna Creek to end of GP property line was surveyed by GP for 108,731ft.
- Georgia-Pacific. 1996. Stream Inventory Report - Laguna Creek.
- Georgia-Pacific. 1996. Stream Inventory Report - Railroad Gulch.
- Georgia-Pacific. 1996. Stream Inventory Report - East Branch Little North Fork Big River.
- Georgia-Pacific. 1996. Stream Inventory Report - Two Log Creek.
- Georgia-Pacific. 1996. Stream Inventory Report - Hatch Gulch.
- Graham Matthews & Associates (GMA). 2001a. Sediment Source Analysis for the Big River Watershed, Mendocino County, California. Weaverville, California.
- GMA. 2001b. Sediment source analysis and preliminary sediment budget for the Albion River Watershed, Mendocino County, California. Graham Matthews & Associates. Prepared for Tetra Tech, Inc., Contract 68-C99-249, Work Assignment #0-34.
- Grantham. 2003. Personal Communication.
- Gregory, K.J., Davis, R.J. 1992. Coarse woody debris in stream channels in relation to river channel management in woodland areas. *Regul. Rivers Res. Manage.* Vol. 7, pp 117– 136.
- Guinness Book of World Records. 2006.
http://www.guinnessworldrecords.com/content_pages/record.asp?recordid=47342
- Hallock, R., G. Warner, et al. 1952." California's Part in a Three-State Salmon Fingerling Marking Program. *California Fish and Game* 38: 301-332.
- Harden, D. 1998. *California Geology*. Upper Saddle River, New Jersey. Prentice Hall, Inc.
- Hilborn, R. and C. J. Walters. 1992. *Quantitative Fisheries Stock Assessment: Choice, Dynamics and Uncertainty*. Chapman and Hall, New York.
- Holman, G. and W. Evans. 1964. Noyo River Stream Clearance Projects, California Department of Fish and Game: 12.
- Huntington, C. W. and Frissell, C. A. 1997. Aquatic conservation and salmon recovery in the north coast basin of Oregon: A crucial role for the Tillamook and Clatsop state forests. Prepared for Oregon Trout. Portland, Oregon.
- Hydrology Subcommittee of the Interagency Advisory Committee on Water Data (HSIACWD). 1981 Guidelines for determining flood frequency. U.S. Geological Survey, Office of Water Data Collection, Reston, VA. Survey Bulletin 17B 183 p. (corrected editorially in March 1982).
- Ice, G.G. 1990. Dissolved oxygen and woody debris: detecting sensitive forest streams. Pages 333-46 in Wilhelms, S.C., and J.S. Gulliver, editors. *Air and water mass transfer*. American Society of Civil Engineers, New York, New York.
- Ice, G. G. 2000. The use of active watershed management to achieve or accelerate the accomplishment of watershed goals.

- Jackson, W. 1975. "Big River Was Dammed." Milles Print Company, St. Louis, Missouri.
- Jackson, W. 1991. Mendocino City: A Daily Journal - 1852-1938. Mendocino Historical Research, Inc.
- Jameson, M. 2002. Harvest History in Big River.
- Jones, W. 1986. Salmon Carcass Surveys: North Fork Big River, Little North Fork Big River, and the East Branch Little North Fork Big River, Mendocino County, California. California Department of Fish and Game Memorandum.
- Jones, W. 2000 (Draft). California Coastal Salmon and Steelhead Current Stream Habitat Distribution Table, National Marine Fisheries Service: 127.
- Karr, J. and Dudley D. 1981. Ecological perspective on water quality goals. *Environmental Management* 5:55-68.
- Kauffman, J., R. Besscha, et al. 1997. "An ecological perspective of riparian and stream restoration in the western United States." *Fisheries* 22(5): 12-24.
- Kight, T. and R. Waldman. 2003. Effects of Nest Box Installation on Wood Duck Brood in Big River Watershed. Mendocino High School, School of Natural Resources, Mendocino, California. 9p.
- Kilbourne, R. 1982. Geology and geomorphic features related to landsliding. Glenblair NE (Northspur) 7.5' quadrangle. California Division of Mines and Geology.
- Kilbourne, R. 1983. Geology and geomorphic features related to landsliding. Mendocino 7.5' quadrangle. California Division of Mines and Geology.
- Kilbourne, R. 1983. Geology and geomorphic features related to landsliding. Glenblair SE (Comptche) 7.5' quadrangle. California Division of Mines and Geology.
- Kilbourne, R. 1984. Geology and geomorphic features related to landsliding. Willits NW (Burbeck) 7.5' quadrangle. California Division of Mines and Geology.
- Kilbourne, R. 1984. Geology and geomorphic features related to landsliding. Willits SW (Greenough Ridge) 7.5' quadrangle. California Division of Mines and Geology.
- Kilbourne, R. 1984. Geology and geomorphic features related to landsliding, Boonville NW (Bailey Ridge) 7.5' quadrangle. California Division of Mines and Geology.
- Kilbourne, R. 1986. "Geology and slope stability of the Fort Bragg area, Mendocino County, California." *California Geology* 39(3): 56-67.
- Kilbourne, R. and A. Mata-Sol. 1983. Geology and geomorphic features related to landsliding. Glenblair SW (Mathison Peak) 7.5' quadrangle. California Division of Mines and Geology.
- Kitsap County. 2000. Kitsap Peninsula Salmonid Refugia Study. Port Orchard, Washington.
- Knopp, C. 1993. Testing Indices for Cold Water Fish Habitat, Final Report for the North Coast Regional Water Quality Control Board, California Regional Water Quality Control Board, Santa Rosa, California.
- Koehler, R. D., K. I. Kelson, and G. Matthews. 2001. Sediment storage and transport in the South Fork Noyo River, Jackson State Demonstration Forest. Unpublished Report Submitted To California Department of Forestry and Fire Protection by William Lettis & Associates, Inc. and Graham Matthews & Associates.
- Kroeber, A. 1925. Handbook of the Indians of California. Bureau of America Ethnology Bulletin.
- Leider, S.A. 1989. Increased straying by adult steelhead trout, *Salmo gairdneri*, following the 1980 eruption of Mt. St. Helens. *Environmental Biology of Fishes* 24: 219-229.
- Levins, R. 1969. "Some demographic and genetic consequences of environmental heterogeneity for biological control." *Bulletin of the Entomological Society of America* 15: 237-240.
- Li, H.W., et. al. 1995. *American Fisheries Society Symposium* 17:371-380.
- Lisle, T. E. 1981a. Recovery of aggraded stream channels at gauging stations in northern California and southern Oregon. Pages 189-211 in T. R. H. Davies and A. J. Pearce, eds. *Erosion and Sediment*

- Transport in Pacific Rim Steeplands, Proceedings of the Christchurch Symposium, 25-31 January 1981. International Association of Hydrology Science, Christchurch, New Zealand. Publication Number 132.
- Lisle, T. E. 1981b. Channel recovery from recent large floods in north coastal California: rates and processes. Pages 153-160, in: R. N. Coates (ed.), Proceedings, Symposium on Watershed Rehabilitation in Redwood National Park and Other Pacific Coastal Areas; 25-28 August 1981. Arcata, California. Center for Natural Resources Studies of JMI, Inc, Sacramento, California.
- Lisle, T. E., and M. B. Napolitano. 1998. Effects of recent logging on the main channel of North Fork Caspar Creek. United States Department of Agriculture Forest Service General Technical Report PSW-GTR-168.
- Maahs, B. 1978. Mendocino Coast Restoration Association, Salmon Unlimited.
- Maahs, M. 1999. Fort Bragg's World's Biggest Salmon BBQ: a brief history of the Salmon Restoration Association. Fort Bragg, California.
- Maahs, M. and S. Cannata. 1998. The Albion River Estuary, its history, water quality and use by salmonids, other fish and wildlife species. Humboldt County Resource Conservation District and Coastal Land Trust, Humboldt County, California.
- Madej, M. A. 1982. Sediment transport and channel changes in an aggrading stream in the Puget Lowland, Washington. Pages 5-23 in F. J. Swanson, R. J. Janda, T. Dunne, and D. N. Swanson, editors. Sediment budgets and routing in forested drainage basins. US Department of Agriculture Forest Service Pacific Northwest Research Station, General Technical Report PNW-141.
- Madej, M.A. 1999. What can thalweg profiles tell us? *Watershed Management Council Networker* 8(4): 14-18.
- Madej, M. A. and V. Ozaki. 1996. Channel response to sediment wave propagation and movement, Redwood Creek, California. *Earth Surface Processes and Landforms* 21: 911-927.
- Marcus, L. and S. Reneau. 1981. Historic Sedimentation in an Estuary: Salt Marsh Succession and Change, Big River Estuary, Mendocino County, California. *Watershed Rehabilitation in Redwood National Park and Other Pacific Coastal Areas*, Proceedings of a symposium held August 24-28, 1981. R. Coates, the Center for Natural Resource Studies, JMI, Inc.: 246-257.
- Marshack, J. 2000. A Compilation of Water Quality Goals, California Environmental Protection Agency, Regional Water Quality Control Board Central Valley Region.
- McBain and Trush. 2000. Spawning gravel composition and permeability within the Garcia River watershed, California. Final Report. Prepared for Mendocino County Resource Conservation District, Mendocino, California. 32 pp. without appendices.
- McElhany, P., M. Ruckelshaus, et al. 2000. Viable salmonid populations and recovery of evolutionarily significant units, National Oceanic and Atmospheric Administration.
- Meehan, W. R., Ed. 1991. Influences of forest and rangeland management on salmonids and their habitats, American Fisheries Society.
- Mendocino Coastal Streams Subcommittee of the Advisory Committee on Salmon and Steelhead Trout. 1986. Findings of the Mendocino Coastal Streams Subcommittee of the Advisory Committee on Salmon and Steelhead Trout.
- Mendocino County Department of Planning and Building Services. 1992a. Notice of Public Hearing and Availability of Draft Negative Declaration for Public Review, Ukiah, California.
- Mendocino County Department of Planning and Building Services. 1992b. Staff Report for Use Permit #U 46-91.
- Mendocino County Department of Transportation. 2001. Ten Mile Road Waterborne Sediment Reduction. Mendocino County, California.
- Mendocino Redwood Company (MRC). 2003. Watershed Analysis for Mendocino Redwood Company's Ownership in the Big River Watershed.

- Montgomery D.R., J.M. Buffington, R.D. Schmidt. 1995. Pool spacing in forest channels. *Water Resources Research*, 31: 1097-1104.
- Moyle, P. and G. Sato. 1991. On the design of preserves to protect native fishes. *Battle against extinction: native fish management in the American West*. W. Minkley and J. Deacon. University of Arizona Press, Tucson, Arizona. Pp 155-169.
- Moyle, P. and R. Yoshiyama. 1992. *Fishes, Aquatic Diversity Management Areas, and Endangered Species: A Plan to Protect California's Native Aquatic Biota*. California Policy Seminar, University of California, Davis California.
- Murphy, M. L., W.R. Meehan, Ed. 1991. Chapter 2: Stream ecosystems. *Influences of Forest and Range Management*. American Fisheries Society, Bethesda, MD.
- National Marine Fisheries Service (NMFS). 2001. Status Review Update for Coho Salmon (*Oncorhynchus kisutch*) from the Central California Coast and the California Portion of Southern Oregon/Northern California Coasts Evolutionarily Significant Units. National Marine Fisheries Service, Southwest Fisheries Center, Santa Cruz, California.
- National Park Service (NPS). 2004. *Nationwide Rivers Inventory*.
- Nicholas, J., and D. Hankin. 1988. Chinook salmon populations in Oregon coastal river basins: description of life histories and assessment of recent trends in run strengths. Oregon Department of Fish and Wildlife Research and Development Section, Corvallis. Oregon. 359p. .
- Nielsen, J., M. Maahs, et al. 1991. Anadromous Salmonid Resources of Mendocino Coastal and Inland Rivers 1989-1990. Salmon Trollers Marketing Association, Mendocino County, California under contract with the California Department of Fish and Game: 98.
- North Coast Regional Water Quality Control Board (NCRWQCB). 2000. Review of Russian River Water Quality Objectives for Protection of Salmonid Species Listed Under the Federal Endangered Species Act: 80.
- North Coast Regional Water Quality Control Board (NCRWQCB). 2001. Water Quality Control Plan for the North Coast Region. Amended 2001.
- Panel on the Ecological Integrity of Canada's National Parks. 2004. Report of the Panel on the Ecological Integrity of Canada's National Parks. http://www.pc.gc.ca/docs/pc/rpts/ie-ei/report-rapport_1_E.asp
- Perry, L. 1974. Progress Report: Fisheries Improvement Study, Big River, Mendocino County (Eureka Division, North Coast Project, California). US Fish and Wildlife Service, Portland, Oregon.
- Quinn, P. F., Beven, K., Chevallier, P., et al. 1991. The Prediction of Hillslope Flow Paths for Distributed Hydrological Modeling Using Digital Terrain Models. *Hydrological Processes*, Vol. 5, 59-79.
- Reeves, G. H., L. E. Benda, et al. 1995. A Disturbance-Based Ecosystem Approach to Maintaining and Restoring Freshwater Habitats of Evolutionarily Significant Units of Anadromous Salmonids in the Pacific Northwest. *Evolution and the Aquatic Ecosystem Symposium*. American Fisheries Society, Monterey, California.
- Reice, S. R. 1994. Nonequilibrium determinants of biological community structure. *American Scientist* 82: 424-435.
- Reid, J. 2002. Land Acquisition Evaluation - Big River Acquisition Project, For California Department of Fish and Game by Mendocino Land Trust.
- Reimers, P. 1973. The length of residence of juvenile Chinook salmon in the Sixes River, Oregon. *Fish Commission of Oregon Research Reports* 4(2): 1- 43.
- Reneau, S. 1981. Geography and Geology. *Big River: the Natural History of and Endangered Northern California Estuary*. S. Warrick and E. Wilcox. University of California, Santa Cruz, California. Environmental Field Program: 3-45.

- Reynolds, K., P. Cunningham, et al. 1996. "A design framework for a knowledge-based information management system for watershed analysis in the Pacific Northwest United States." *AI Applications* 10: 9-22.
- Reynolds, K. M. 1999. EMDS Users Guide (version 2.0): Knowledge-based decision support for ecological assessment. Knowledge-based decision support for ecological assessment. , U.S. Department of Agriculture. Pacific Northwest Research Station, U. S, Forest Service, Portland, OR. 63p.
- Rosgen, D. 1996. *Applied River Morphology*. Wildland Hydrology, Pagosa Springs, Colorado.
- Salmon Trollers Marketing Association. 1990. *Salmon Trollers Stream Restoration Project Preliminary Report for February 1990 - Louisiana Pacific Property*.
- Scriven, J. 2003. *Big River Watershed Stream Restoration Project Final Report*. Ukiah, California. Prepared for US Fish and Wildlife Service by E Center's Mendocino Fisheries Program.
- Seacat, M., D. Seymour, et al. 1981. *Vegetation. Big River: the Natural History of an Endangered Northern California Estuary*. S. Warrick and E. Wilcox. University of California, Santa Cruz, California, Environmental Field Program. pp 100-161.
- Sedell, J., G. Reeves, et al. 1990. "Role of refugia in recovery from disturbances: modern fragmented and disconnected river systems." *Environmental Management* 14(5): 711-724.
- Sharples, N. 2003. *Little North Fork Big River Coho and Steelhead Carcass/Redd Survey*. Mendocino High School, School of Natural Resources, Mendocino, California. 8p.
- Shirazi, M., W. Seim, et al. 1979. *Characterization of Spawning Gravel and Stream System Evaluation*, US Environmental Protection Agency.
- Short, W. and Spittler. 2002. *GIS Data for the watersheds mapping series, map set 2, Jackson Demonstration State Forest Mendocino County, California*, California Geological Survey. CD 2002-05.
- Sommarstrom, S. 1984. *Mendocino County Salmon and Steelhead Management Plan*. Mendocino County Board of Supervisors, Ukiah, California. 103p.
- Mendocino High School of Natural Resources (SONAR). 2001. *Carcass and Redd Survey of Little North Fork Big River*.
- SONAR. 2001. *Carcass and Redd Survey of Railroad Creek*.
- SONAR. 2001. *Carcass and Redd Survey of West Chamberlain Creek*.
- SONAR. 2001. *Carcass and Redd Survey of Chamberlain Creek*.
- SONAR. 2002. *SONAR Big River Estuary Study Mud Flat #1 Plant Species List*.
- SONAR. 2002. *Carcass and Redd Survey of Little North Fork Big River*.
- SONAR. 2003. *Carcass and Redd Survey of Little North Fork Big River*.
- SONAR. 2002. *Wood Duck Box Locations and GPS Coordinates*. Mendocino, California.
- Sullivan, K. D.J. Martin, R.D. Cardwell, J.E. Toll, and S. Duke. 2000. *An analysis of the effects of temperature on salmonids of the Pacific Northwest with implications for selecting temperature criteria*. Sustainable Ecosystem Institute, Portland Oregon. (DRAFT).
- Swanston, D. N. 1991. *Natural Processes*. Pages 139-179 in W. R. Meehan, editor. *Influences of forest and rangeland management on salmonid fisheries and their habitats*. American Fisheries Society Special Publication 19. Bethesda, Maryland.
- Taylor, R. N. 2001. *Final Report: Coastal Mendocino County Culvert Inventory and Fish Passage Evaluation*. CDFG, McKinleyville, CA. 43p.
- United States Bureau of Reclamation (USBR). 1974. *Fishery Improvement Study, Eureka Division, North Coast Project, California*. Department of the Interior, Bureau of Reclamation, Sacramento, California. 46p.

- United States Environmental Protection Agency (EPA). 2001. Big River Total Maximum Daily Load for Sediment (TMDL).
- Ven te Chow. 1964. Handbook of applied hydrology. McGraw-Hill, New York.
- WAC Corporation. 1984. Black and white aerial photos, Flight 84C, Flight Line 15, Frames 44-49, 151-162, 213-222; Flight Line 18, Frames 6-8, 57-63, 113-117, 170-176, 227-236; Flight Line 20, Frames 106-117; scale 1:31,680.
- WAC Corporation. 2000. Black and white aerial photos, Flight WAC-00-CA, Flight Line 2, Frames 3-13, 29-41, 56-65, 81-92, 108-113, 210-212; Flight Line 3, Frames 208-210; Flight Line 10, Frames 82-91, 139-142, 195-202; Flight Line 11, Frames 6-12, 49-55, 83-90, 111-119, 137-147; scale 1:24,000.
- Washington Forest Practice Board. 1997. Board Manual: Standard Methodology for Conducting Watershed Analysis. Version 4.0. Washington Forest Practice Board, Olympia, WA.
- Warrick, S. and E. Wilcox, Eds. 1981. Big River: the Natural History of an Endangered Northern California Estuary. Environmental Field Program, University of California Santa Cruz. Santa Cruz, California.
- Wright, C. 2004. Personal communication.
- Wurm, T. 1986. Mallets on the Mendocino Coast: Casper Lumber Company, Mallets on the Mendocino Coast: railroads and steamships. Trans-Anglo Books, Glendale, California.
- Wynn, R. 1989-1990. "Bridges, Huckleberries, and Robin Stew: The Depression and the New Deal in Mendocino County." Mendocino Historical Society Review XIV. Mendocino, California.

Spatial Data Availability, Catalog, Standards and Analyses

Data Availability

For GIS data associated with this report please contact the individual agencies listed below:

CaSIL – California Spatial Information Library (formerly Teale – Stephen P. Teale data center, State of California): <http://gis.ca.gov/>

CDF – California Department of Forestry and Fire Protection: www.fire.ca.gov

CGS – Department of Conservation, California Geological Survey: www.consrv.ca.gov/CGS

CWMC – California Watershed Mapping Committee: <http://www.ca.nrcs.usda.gov/features/calwater>

CWPAP – Coastal Watershed Planning and Assessment Program: <http://www.coastalwatersheds.ca.gov>

CDFG – California Department of Fish and Game: <http://www.dfg.ca.gov>

DOD – Department of Defense: <http://www.defenselink.mil/>

DWR – California Department of Water Resources: <http://www.water.ca.gov/>

FRAP – Forest Resource Assessment Program: <http://frap.cdf.ca.gov/>

NCRWQCB – North Coast Regional Water Quality Control Board: <http://www.swrcb.ca.gov/rwqcb1/>

RNSP – Redwood National and State Parks: <http://www.nps.gov/redw>

SSRRC – Sotoyome-Santa Rosa Resource Conservation District: <http://sotoyomercd.org/>

USDA – United States Department of Agriculture: <http://www.usda.gov/wps/portal/usdahome>

USGS – United States Geological Survey: <http://www.usgs.gov/>

Spatial and Geographic Information Systems (GIS) Data Standards and Analyses

Data records were collected for synthesis and analysis purposes and most of these data were either created in a spatial context or converted to a spatial format. Effective use of these data between the five partner departments required establishing standards for data format, storage, management and dissemination. Early in the assessment process, we held a series of meetings designed to gain consensus on a common format for the often widely disparate data systems within each department. Our objective was to establish standards that could be easily used by each department, that were most useful and powerful for selected analysis, and would be most compatible with standards used by potential private and public sector stakeholders.

As a result, we agreed that spatial data and base information disseminated to the public through the program would be in the following format (See data catalog for a complete description of data sources and scale):

Data form: standard database format usually associated with a GIS shapefile[®] (ESRI) or coverage. Data were organized by watershed and distributed among watershed synthesis teams. Electronic images were retained in their current format.

Spatial Data Projection: spatial data were projected from their native format to both Teale Albers, North American Datum (NAD) 1927 and Universal Transverse Mercator (UTM), Zone 10, NAD 1983. Both formats were used in data analysis and synthesis.

Scale: most data were created and analyzed at 1:24000 scale to 1) match the minimum analysis scale for planning watersheds, and 2) coincide with base information (e.g., stream networks) on USGS quadrangle maps (used as Digital Raster Graphics [DRG]).

Data Sources: data were obtained from a variety of sources including spatial data libraries with partner departments or were created by manually digitizing from 1:24000 DRG.

The metadata available for each spatial data set contain a complete description of how data were collected and attributed. Spatial data sets that formed the foundation of most analysis included the 1:24000 hydrography and the 10-meter scale Digital Elevation Models (DEM). Hydrography data were created by manually digitizing from a series of 1:24000 DRG then attributing with direction, routing, and distance information using a dynamic segmentation process (see

<http://support.esri.com/index.cfm?fa=knowledgebase.whitepapers.viewPaper&PID=43&MetaID=294> for more

information). The resulting routed hydrography allowed for precise alignment and display of stream habitat data and other information along the stream network. The DEM was created from base contour data obtained from the USGS for the entire assessment region.

Source spatial data were often clipped to watershed, planning watershed, and subbasin units prior to use in analysis. Analysis often included creation of summary tables, tabulating areas, intersecting data based on selected attributes, or creation of derivative data based on analytical criteria. For more information regarding the approach to analysis and basis for selected analytical methods, see the integrated analyses section.